

'Where every day is an opportunity to shine.'

Is my child too ill for school?

It can be tricky deciding whether or not to keep your child off school when they're unwell.

There are government guidelines for schools about managing specific infectious diseases at GOV.UK. These say when children should be kept off school and when they shouldn't.

If you do keep your child at home, it's important to phone the school or nursery on the first day. Let them know that your child won't be in and give them the reason. If your child is well enough to go to school but has an infection that could be passed on, such as a cold sore or headlice, let their teacher know.

For other illnesses follow this advice:

Coughs and colds

It's fine to send your child to school with a minor cough or common cold. But if they have a fever, keep them off school until the fever goes.

Encourage your child to throw away any used tissues and to wash their hands regularly.

High temperature

If your child has a high temperature, keep them off school until it goes away.

Chickenpox

If your child has chickenpox, keep them off school until all the spots have crusted over. Thisisusuallyabout5 days after the spots first appeared

Cold sores

There's no need to keep your child off school if they have a cold sore. Encourage them not to touch the blister or kiss anyone while they have the cold sore, or to share things like cups and towels.

Conjunctivitis

You don't need to keep your child away from school if they have conjunctivitis. Do get advice from your pharmacist. Encourage your child not to rub their eyes and to wash their hands regularly.

COVID-19

If your child has mild symptoms, such as a runny nose, sore throat, or slight cough, and feels well enough, they can go to school.

Your child should try to stay at home and avoid contact with other people if they have symptoms of COVID-19 and they either:

have a high temperature do not feel well enough to go to school do their normal activities.

What to do if your child has tested positive:

Your child is no longer required to do a COVID-19 rapid lateral flow test if they have symptoms.

But if your child has tested positive for COVID-19, they should try to stay at home and avoid contact with other people for 3 days after the day they took the test.

Ear infection

If your child has an ear infection and a high temperature or severe earache, keep them off school until they're feeling better or their high temperature goes away.

Hand, foot and mouth disease

If your child has hand, foot and mouth disease but seems well enough to go to school, there's no need to keep them off.

Encourage your child to throw away any used tissues straight away and to wash their hands regularly.

Head lice and nits

There's no need to keep your child off school if they have headlice. Do try to treat them and get them.

straight back into school. You can treat headlice and nits without seeing a GP.

Impetigo

If your child has impetigo, they'll need treatment from a GP, often with antibiotics. Keep them off school until all the sores have crusted over and healed, orfor48 hours after they start antibiotic treatment.

Encourage your child to wash their hands regularly and not to share things like towels and cups with other children at school.

Ringworm

If your child has ringworm, see your pharmacist unless it's on the scalp, in which case you should see a GP. It is fine for your child to go to school once they have started treatment.

Scarlet fever

If your child has scarlet fever, they'll need treatment with antibiotics from a GP.
Otherwise, they'll be infectiousfor2 to3 weeks.

Your child can return to school 24 hours after starting antibiotics.

Slapped cheek syndrome (fifth disease)

You don't need to keep your child off school if they have slapped cheek syndrome because, once the rash appears, they're no longer infectious.

But let the school or teacher know if you think your child has slapped cheek syndrome.

Sore throat

You can still send your child to school if they have a sore throat. But if they also have a high temperature, they should stay at home until it goes away.

A sore throat and high temperature can be symptoms of tonsillitis.

Threadworms

You don't need to keep your child off school if they have threadworms. Speak to your pharmacist, who can recommend a treatment.

Vomiting and diarrhoea

Children with diarrhoea or vomiting should stay away from school until they have not been sick or had diarrhoea for at least 2 days (48 hours)

